

AD-A049 386

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

RATIONAL BOUNDS FOR THE T-TAIL AREA WITH AN APPLICATION TO BONF--ETC(U)

NOV 77 A P SOMS

DAAG29-75-C-0024

UNCLASSIFIED

MRC-TSR-1807

NL

| OF |

ADA049386



END  
DATE  
FILMED  
3-78  
DDC

AD A049386

AD NO. FILE COPY

MRC Technical Summary Report #1807


RATIONAL BOUNDS FOR THE t-TAIL AREA  
WITH AN APPLICATION TO BONFERRONI  
t-PERCENTILES

Andrew P. Soms

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

November 1977

Received October 7, 1977



Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

and

University of Wisconsin-Milwaukee  
Graduate School

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

Rational Bounds for the t-Tail Area  
With an Application to Bonferroni t-Percentiles

Andrew P. Soms

Technical Summary Report #1807  
November 1977

Abstract


The bounds of Boyd (1959) for the normal distribution are extended to the t-distribution. It is shown how these bounds can be used to calculate the Bonferroni descriptive level as well as Bonferroni t-percentiles. The adequacy of the approximation is discussed and numerical examples provided.

Key words: Bonferroni percentiles, bounds, descriptive Bonferroni level, t-tail area, t-distribution.

AMS(MOS) Subject Classification: 60E05

Work Unit No. 4 - Probability, Statistics and Combinatorics

ACCESSION No.	
NTIS	White Series <input checked="" type="checkbox"/>
DDC	Red Series <input type="checkbox"/>
UNANNOUNCED <input type="checkbox"/>	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. AND/OR SPECIAL
A	



Sponsored by the United States Army under Contract No. DAAG29-75-C-0024  
and the University of Wisconsin-Milwaukee Graduate School.

see 1473

## SIGNIFICANCE AND EXPLANATION

In experimental work it is sometimes desired to make  $n$  different inferences from a given data set. Associated with the data is a hypothesis called the overall null hypothesis which is to be tested, and under the assumption that the overall null hypothesis is true it is desired to have a preassigned high probability  $1-\alpha$  of having all the individual inferences be correct. If the proper assumptions are made concerning the statistical distributions involved, this requires finding the abscissa  $t_{\alpha/2n}$  for a specified t-distribution having area  $\alpha/(2n)$  to the right of it. Such abscissas are called Bonferroni t-percentiles. Since statistical tables only give abscissas corresponding to  $\alpha$ , it is sometimes troublesome to find  $t_{\alpha/2n}$ . Whereas previous work of the author is applicable if  $\alpha/(2n)$  is small, the present paper gives a method that works well for any  $\alpha/(2n)$  between .0 and .5 and is readily implemented on a computer.



## Rational Bounds for the t-Tail Area

### With an Application to Bonferroni t-Percentiles

Andrew P. Soms

The purpose of this paper is to complete the extension to the t-distribution of results known for the normal (see, e.g., Johnson and Kotz 1970, Chapter 33). In Soms (1976) an asymptotic expansion for the tail area of the t-distribution was given good for small areas and in Soms (1977) some improved bounds were given, none of which, however, had simultaneously the right limiting behavior as  $x \rightarrow 0$  or  $x \rightarrow \infty$ . In this paper lower and upper bounds are given with this property, the accuracy of the approximation is studied, and the results applied to the calculation of Bonferroni t-percentile and descriptive levels.

#### 2. Extensions of Boyd's Results

Boyd (1959) showed that if  $\phi(x) = (2\pi)^{-1/2} e^{-x^2/2}$ ,  $\bar{F}_k(x) = \int_x^\infty \phi(t) dt$ , and  $R_x = \bar{F}_k(x)/\phi(x)$ ,  $x > 0$ , then

$$p(x, \gamma_{\min}) < R_x < p(x, \gamma_{\max}),$$

where  $p(x, \gamma) = (\gamma+1)/[(x^2 + \frac{2}{\pi}(\gamma+1)^2)^{1/2} + \gamma x]$ ,  $\gamma_{\max} = 2/(\pi-2)$ ,  $\gamma_{\min} = \pi-1$ , and the bounds are the best possible in the class  $\{p(x, \gamma), \gamma > -1\}$ . This is also discussed in Johnson and Kotz (1970, Chapter 33). Here we will obtain analogous results for the t-distribution.

Let

$$f_k(t) = c_k (1+t^2/k)^{-(k+1)/2}, \quad c_k = \frac{\Gamma((k+1)/2)}{\Gamma(k/2)(\pi k)^{1/2}},$$

k an integer  $\geq 1$ , and

$$\bar{F}_k(x) = 1 - F_k(x) = \int_x^\infty f_k(t) dt,$$

for  $x > 0$  here and throughout the paper. Also let

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024 and the University of Wisconsin-Milwaukee Graduate School.

$$R_x = \bar{F}_k(x) / ((1+x^2/k)f_k(x)) .$$

was shown in Soms (1976) that

$$1/x - k/((k+2)x^3) < R_x < 1/x$$

and

$$R_x = 1/x - k/((k+2)x^3) + O(x^{-5}) . \quad (2.1)$$

In Soms (1977) it was shown that

$$\left[ \frac{(k-1)x}{2k} + \left( 1 + \left( \frac{(k+1)x}{2k} \right)^2 \right)^{1/2} \right]^{-1} < R_x < \left[ \frac{(3k-1)x}{4k} + \left( \frac{k-1}{2k} + \left( \frac{(k+1)x}{4k} \right)^2 \right)^{1/2} \right]^{-1} , \quad (2.2)$$

where the lower bound is valid for  $k \geq 1$  and the upper for  $k \geq 2$ . It can be seen that these bounds do not have the right limiting behavior as  $x \rightarrow 0$ . Motivated by (2.2), consider the problem of finding best upper and lower bounds of the form

$$\frac{\alpha}{(x^2 + \beta)^{1/2} + \gamma x} .$$

Since  $\lim_{x \rightarrow 0} R_x = 1/2c_k$  and  $R_x \sim 1/x$  (as  $x \rightarrow \infty$ ), one must have

$$\alpha = 1 + \gamma \quad \text{and} \quad \alpha/\beta^{1/2} = 1/2c_k .$$

Thus  $\beta = 4c_k^2(1+\gamma)^2$  and the approximating functions  $p(x, \gamma)$  must be of the form

$$p(x, \gamma) = \frac{1+\gamma}{(x^2 + 4c_k^2(1+\gamma)^2)^{1/2} + \gamma x} .$$

For small  $x$ ,

$$R_x = 1/2c_k - x + O(x^2)$$

and

$$p(x, \gamma) = 1/2c_k - \gamma x / (4c_k^2(1+\gamma)) + O(x^2) .$$

For large  $x$ , from (2.1),

$$R_x = 1/x - k/((k+2)x^3) + O(x^{-5})$$

and

$$p(x, \gamma) = 1/x - 2c_k^2(1+\gamma)/x^3 + O(x^{-5}) .$$

So if  $p(x, \gamma)$  is to maximize  $R_x$  it is necessary that

$$\frac{\gamma}{4c_k^2(\gamma+1)} \leq 1 \quad \text{and} \quad \frac{k}{k+2} \geq 2c_k^2(1+\gamma) ,$$

or

$$-1 < \gamma \leq \frac{4c_k^2}{1-4c_k^2} \quad \text{and} \quad -1 < \gamma \leq \frac{k}{2(k+2)c_k^2} - 1 ,$$

or

$$-1 < \gamma \leq \min \left( \frac{4c_k^2}{1-4c_k^2} , \frac{k}{2(k+2)c_k^2} - 1 \right) . \quad (2.3)$$

Note that from Lemmas A2 and A3 and the direct evaluation of  $\frac{k}{2(k+2)c_k^2} - 1$

for  $k = 1$  the interval in (2.3) is non-empty. Similarly, if  $p(x, \gamma)$  is to minimize  $R_x$ , it is necessary that

$$\frac{\gamma}{4c_k^2(\gamma+1)} \geq 1 \quad \text{and} \quad \frac{k}{k+2} \leq 2c_k^2(\gamma+1) ,$$

or

$$\gamma \geq \max \left( \frac{4c_k^2}{1-4c_k^2} , \frac{k}{2(k+2)c_k^2} - 1 \right) .$$

Now

$$\frac{4c_k^2}{1-4c_k^2} \geq \frac{k}{2(k+2)c_k^2} - 1$$

is equivalent to

$$c_k^2 \geq k/(6k+4).$$

Hence from Lemma A4, if  $k \geq 3$ , then for maximizing it is necessary that

$$\gamma \leq 4c_k^2/(1-4c_k^2),$$

and for minimizing

$$\gamma \geq \frac{k}{2(k+2)c_k^2} - 1.$$

If  $k = 2$ , then  $4c_k^2/(1-4c_k^2) = k/(2(k+2)c_k^2) - 1$ , and it is verified by differentiation that (here and throughout "'" will denote differentiation)

$$- \left[ \left( 1 + \frac{x^2}{k} \right) f_k(x) p(x, \gamma) \right]' = f_k(x),$$

and hence in this case  $R_x = p(x, \gamma)$ , where  $\gamma = 4c_k^2/(1-4c_k^2) = k/(2(k+2)c_k^2) - 1$ .

Therefore for  $k = 2$  the approximating problem has been solved in the best possible way and this case will not be considered further. If  $k = 1$ , then for maximizing it is necessary that

$$-1 < \gamma \leq \frac{k}{2(k+2)c_k^2} - 1,$$

and for minimizing

$$\gamma \geq 4c_k^2/(1-4c_k^2),$$

again using Lemma A4.



Using the above as motivation, the first result is

Theorem 2.1: Let  $\gamma = 4c_k^2/(1-4c_k^2)$ . Then for  $k \geq 3$ ,  $R_x < p(x, \gamma)$  and for  $k = 1$ ,  $R_x > p(x, \gamma)$ .

Proof: Note that  $R'_x = ((\frac{k-1}{k})x R - 1)/(1+x^2/k)$  and  $R''_x = ((\frac{k-1}{k})(\frac{(k-2)x^2}{k} + 1)R_x - \frac{(k-3)x}{k})/(1+x^2/k)$ . Let  $f(x) = p(x, \gamma) - R_x$ . Then  $f(0) = f'(0) = 0$  and

$$\begin{aligned} f''(0) &= \frac{-1+2\gamma^2}{8c_k^3(1+\gamma)^2} - \frac{k-1}{2kc_k} \\ &= \frac{16kc_k^4 + (4k+4)c_k^2 - k}{8c_k^3(1+\gamma)^2 k} . \end{aligned}$$

It is verified directly that  $f''(0) < 0$  for  $k = 1$ . For  $k \geq 3$ ,  $f''(0) > 0$  is equivalent to

$$16kc_k^4 + (4k+4)c_k^2 > k , \quad (2.4)$$

which will certainly be true if

$$16c_k^4 + 4c_k^2 > 1 . \quad (2.5)$$

In Lemma A1 it is shown that the even and odd  $c_k$ 's form an increasing sequence. It follows by direct verification that (2.4) is true for  $3 \leq k \leq 16$  and that (2.5) is true for  $k = 17$  and  $k = 18$  and hence (2.4) is true for  $k \geq 3$ . So for  $k \geq 3$ ,  $f''(0) > 0$ . Consider

$$h(x) = \frac{k-1}{k} xf/(1+x^2/k) - f' .$$

By some algebra, it follows that

$$\begin{aligned} &(1+x^2/k) \left[ (x^2+4c_k^2(1+\gamma)^2)^{\frac{1}{2}} + \gamma x \right]^2 (x^2+4c_k^2(1+\gamma)^2)^{\frac{1}{2}} (1-4c_k^2)^3 h(x) \\ &= x \left[ 4c_k^2(8c_k^2-1) \left( \frac{x}{(x^2+4c_k^2/(1-4c_k^2))^2 + x} \right) - \left( -1 + 4 \frac{k+1}{k} c_k^2 + 16c_k^4 \right) \right] . \quad (2.6) \end{aligned}$$

By Lemma A2 and A3,  $1-4c_k^2 > 0$  for  $k \geq 1$ , and  $8c_k^2 - 1 > 0$  for  $k \geq 3$  and  $8c_k^2 - 1 < 0$  for  $k = 1$ . Also, as  $x \rightarrow \infty$ ,

$$4c_k^2(8c_k^2-1) \frac{x}{(x^2+4c_k^2/(1-4c_k^2)^2+x)} + 2c_k^2(8c_k^2-1), \quad (2.7)$$

and the convergence is monotone. For  $k \geq 3$ ,

$$2c_k^2(8c_k^2-1) > -1 + 4 \frac{k+1}{k} c_k^2 + 16c_k^4 \quad (2.8)$$

and for  $k = 1$ ,

$$2c_k^2(8c_k^2-1) < -1 + 4 \frac{k+1}{k} c_k^2 + 16c_k^4, \quad (2.9)$$

since (2.8) and (2.9) are equivalent to  $c_k^2 < \frac{k}{6k+4}$  and  $c_k^2 > \frac{k}{6k+4}$ , respectively, and this is true by Lemma A5. Since the constant term in the parentheses of the right-hand side of (2.6) has the same sign as  $f''(0)$ , it follows from (2.7) and (2.8) that for  $k \geq 3$

$$h(x) < 0, \quad x < x_0, \quad (2.10)$$

$$h(x) = 0, \quad x = x_0,$$

and

$$h(x) > 0, \quad x > x_0. \quad (2.11)$$

Suppose now that  $f(x) \leq 0$  for some  $x \leq x_0$ . Then since  $f(x)$  is increasing at the origin, it must have a maximum in  $(0, x_0)$  for which  $f > 0$  and  $f' = 0$ , contradicting (2.10). If  $f(x) \leq 0$  for some  $x > x_0$ , then  $f$  must have a minimum ( $\lim_{x \rightarrow \infty} f(x) = 0$ ) for which  $f \leq 0$  and  $f' = 0$ , contradicting (2.11). So  $R_x < p(x, \gamma)$  for  $k \geq 3$ .

Consider now  $k = 1$ . By (2.7), (2.9), and the fact that  $f''(0) < 0$  for  $k = 1$ ,

$$h(x) > 0, \quad x < x_0, \quad (2.12)$$

$$h(x) = 0, \quad x = x_0,$$

and

$$h(x) < 0, \quad x > x_0. \quad (2.13)$$

Suppose that  $f(x) \geq 0$  for some  $x \leq x_0$ . Then, since  $f$  is decreasing at the origin there must be an  $x < x_0$  for which  $f < 0$  and  $f' = 0$ , contradicting (2.12). If  $f(x) \geq 0$  for  $x > x_0$ , there must be an  $x > x_0$  for which  $f \geq 0$  and  $f' = 0$ , contradicting (2.13). Hence for  $k = 1$ ,  $R_x > p(x, \gamma)$ , completing the proof.

Consider now the problem of minimizing  $R_x$  for  $k \geq 3$  and maximizing for  $k = 1$ . Using similar reasoning to that used to arrive at the statement of Theorem 2.1, we have

Theorem 2.2: Let  $\gamma = \frac{k}{2(k+2)c_k^2} - 1$ . Then for  $k \geq 3$ ,  $R_x > p(x, \gamma)$  and for

$$k = 1, R_x < p(x, \gamma).$$

Proof: Let  $g(x) = R_x - p(x, \gamma)$ . Then  $g(0) = 0$  and

$$\begin{aligned} g'(0) &= -1 + \left( \frac{k}{2(k+2)c_k^2} - 1 \right) \sqrt{4c_k^2 \frac{k}{2(k+2)c_k^2}} \\ &= \frac{(-6k-4)c_k^2 + k}{4c_k^2}, \end{aligned}$$

and hence by Lemma A5,  $g'(0) > 0$  for  $k \geq 3$ , and  $g'(0) < 0$  for  $k = 1$ . Consider

$$h(x) = \frac{k-1}{k} xg/(1+x^2/k) - g'.$$

After some algebra,

$$\begin{aligned} h &= \left[ -(x^2 + 4c_k^2(1+\gamma)^2)^{\frac{1}{2}} ((\gamma-1)x^2 + (1+\gamma)(-4c_k^2(1+\gamma) + \gamma)) \right. \\ &\quad \left. + x \left\{ (\gamma-1)x^2 + (1+\gamma)(4c_k^2(1+\gamma)((1+1/k)\gamma - 1 + 1/k) - 1) \right\} \right] \\ &\quad / \left[ ((x^2 + 4c_k^2(1+\gamma)^2)^{\frac{1}{2}} + \gamma x)^2 (x^2 + 4c_k^2(1+\gamma)^2)^{\frac{1}{2}} (1+x^2/k) \right], \end{aligned}$$

or, upon some further algebraic simplification,

$$\begin{aligned}
& ((x^2 + 4c_k^2(1+\gamma)^2)^{\frac{3}{2}} + \gamma x)^2 (x^2 + 4c_k^2(1+\gamma)^2)^{\frac{3}{2}} (1+x^2/k) ((x^2 + 4c_k^2(1+\gamma)^2)^{\frac{3}{2}} + x) h(x) / (1+\gamma)^2 \\
&= -4c_k^2 \left[ x^2(\gamma-1) + (1+\gamma)(-4c_k^2(1+\gamma) + \gamma) \right] \\
&+ x^2 \left[ 4c_k^2((1+1/k)\gamma + 1/k) - 1 \right] + (x^2 + 4c_k^2(1+\gamma)^2)x \\
&\cdot \left[ 4c_k^2((1+1/k)\gamma + 1/k) - 1 \right]. \tag{2.14}
\end{aligned}$$

Note that

$$4c_k^2((1+1/k)\gamma + 1/k) - 1 = 4c_k^2(\gamma-1) - 4c_k^2((1+1/k)\gamma + 1/k) + 1. \tag{2.15}$$

Hence, using (2.15), the right-hand side of (2.14), apart from the constant term, can be written as

$$-cx^2 + cx(x^2 + 4c_k^2(1+\gamma)^2)^{\frac{3}{2}} = \frac{c^2 4c_k^2(1+\gamma)^2}{c[1+(1+4c_k^2(1+\gamma)^2/x^2)^{\frac{3}{2}}]}, \tag{2.16}$$

where  $c = 4c_k^2((1+1/k)\gamma + 1/k) - 1$ . Also,  $-4c_k^2(1+\gamma) + \gamma \geq 0$  is equivalent to  $c_k^2 \leq k/(6k+4)$ , and hence by Lemma A5,  $-4c_k^2(1+\gamma) + \gamma > 0$  for  $k \geq 3$  and  $-4c_k^2(1+\gamma) + \gamma < 0$  for  $k = 1$ . Further,  $4c_k^2((1+1/k)\gamma + 1/k) - 1 > 0$  is equivalent to

$$c_k^2 < k/(4k+8),$$

and since

$$k/(4k+8) > k/(6k+4), \quad k \geq 3,$$

by Lemma A5

$$4c_k^2((1+1/k)\gamma + 1/k) - 1 > 0, \quad k \geq 3.$$

It is verified by direct evaluation that  $4c_k^2((1+1/k)\gamma + 1/k) - 1 < 0$  for  $k = 1$ .



Consider the limit  $2cc_k^2(1+\gamma)^2$  of (2.16) as  $x \rightarrow \infty$ . For  $k \geq 3$ ,

$$2cc_k^2(1+\gamma)^2 > 4c_k^2(1+\gamma)(-4c_k^2(1+\gamma)+\gamma) \quad (2.17)$$

is equivalent to

$$c_k^2 > \frac{(k)(k+4)}{8(k+1)(k+2)} \quad (2.18)$$

(2.18) will certainly be true if

$$c_k^2 > \frac{k+4}{8(k+2)}, \quad (2.19)$$

and it is verified directly that (2.19) holds for  $k = 9$  and  $k = 10$  and hence (2.17) is true for  $k \geq 9$  by Lemma A1. It is also verified directly that (2.17) holds for  $3 \leq k \leq 8$  and that the inequality is reversed for  $k = 1$ .

Hence it follows that for  $k \geq 3$

$$h(x) < 0, \quad x < x_0, \quad (2.20)$$

and

$$\begin{aligned} h(x) &= 0, & x &= x_0, \\ h(x) &> 0, & x &> x_0, \end{aligned} \quad (2.21)$$

and for  $k = 1$

$$h(x) > 0, \quad x < x_0, \quad (2.22)$$

and

$$\begin{aligned} h(x) &= 0, & x &= x_0, \\ h(x) &< 0, & x &> x_0. \end{aligned} \quad (2.23)$$

Consider  $k \geq 3$  first and suppose that  $g(x) \leq 0$  for some  $x \leq x_0$ . Then, since  $g$  is increasing at the origin, there must be an  $x$  in  $(0, x_0)$  such that  $g > 0$  and  $g' = 0$ , which contradicts (2.20). If  $g(x) \leq 0$  for some  $x > x_0$  then there must be an  $x > x_0$  for which  $g \leq 0$  and  $g' = 0$  and this contradicts (2.21). Hence  $R_x > p(x, \gamma)$  for  $k \geq 3$ . If  $k = 1$  and  $g(x) \geq 0$  for some  $x \leq x_0$ , then, since  $g$  is decreasing at the origin, there must be an  $x$

in  $(0, x_0)$  such that  $g(x) < 0$  and  $g'(x) = 0$ , contradicting (2.22). If  $g(x) \geq 0$  for some  $x > x_0$ , then there must be an  $x > x_0$  for which  $g \geq 0$  and  $g' = 0$ , contradicting (2.23) and completing the proof of Theorem 2.2.

We summarize the results of Theorem 2.1 and 2.2 below in an algebraically equivalent form. For  $k \geq 3$ , let  $\gamma_{\max} = 4c_k^2/(1-4c_k^2)$  and

$$\gamma_{\min} = \frac{k}{2(k+2)c_k^2} - 1. \text{ Then}$$

$$\left[ \left( \left( \frac{x}{(1+\gamma_{\min})} \right)^2 + 4c_k^2 \right)^{\frac{1}{2}} + \gamma_{\min} x / (1+\gamma_{\min}) \right]^{-1} < R_x < \left[ \left( \left( \frac{x}{(1+\gamma_{\max})} \right)^2 + 4c_k^2 \right)^{\frac{1}{2}} + \gamma_{\max} x / (1+\gamma_{\max}) \right]^{-1}. \quad (2.24)$$

If  $k = 2$ , then  $\gamma_{\max} = \gamma_{\min} = \gamma$  and

$$R_x = \left[ \left( \left( \frac{x}{(1+\gamma)} \right)^2 + 4c_k^2 \right)^{\frac{1}{2}} + \gamma x / (1+\gamma) \right]^{-1}.$$

For  $k = 1$ , (2.24) holds with the definitions of  $\gamma_{\max}$  and  $\gamma_{\min}$  interchanged.

Since  $\partial p(x, \gamma) / \partial \gamma < 0$  for  $x > 0$ , it follows that for  $k \geq 3$ ,  $\gamma_{\max}$  is the best value of  $\gamma$  for which  $R_x < p(x, \gamma)$  for all  $x$  and  $\gamma_{\min}$  is the best value for which  $R_x > p(x, \gamma)$  for all  $x$ . A similar statement holds for  $k = 1$  with the definitions of  $\gamma_{\max}$  and  $\gamma_{\min}$  interchanged. Equivalently, the bounds given in Theorems 2.1 and 2.2 are uniformly best in the class considered.

The accuracy of the lower and upper bounds are related to the maximum value of the ratio  $p(x, \gamma_{\max}) / p(x, \gamma_{\min})$ . Call the ratio  $r(x)$  and consider the maximum value of  $r(x)$ , which must be attained for some positive  $x$  at which  $r'(x) = 0$ . By differentiation,  $r'(x) = 0$  is equivalent to

$$f(x) = (a_1^2 - a_2^2)x + (1-a_1)(a_1^2 x^2 + 4c_k^2)^{\frac{1}{2}} - (1-a_2)(a_2^2 x^2 + 4c_k^2)^{\frac{1}{2}} = 0, \quad (2.25)$$

where  $a_1 = 2(k+2)c_k^2/k$  and  $a_2 = 1-4c_k^2$  if  $k \geq 3$ , and the definitions of  $a_1$  and  $a_2$  are interchanged if  $k = 1$ . It will now be shown that  $f$  as defined by (2.25) is a monotonically decreasing function,  $f(0) > 0$ ,  $f(\infty) < 0$ , and hence the maximum is the unique solution of (2.25). By differentiation,

$$f'(x) = a_1^2 - a_2^2 + \frac{(1-a_1)a_1^2 x}{(a_1^2 x^2 + 4c_k^2)^{3/2}} - \frac{(1-a_2)a_2^2 x}{(a_2^2 x^2 + 4c_k^2)^{3/2}}. \quad (2.26)$$

Consider  $k \geq 3$ , a similar argument holding for  $k = 1$ .  $a_1 < a_2$  is equivalent to  $c_k^2 < k/(6k+4)$ , true by Lemma A5. Hence  $a_1^2 - a_2^2 < 0$  and

$$f'(x) < a_1^2 - a_2^2 + \frac{a_2^2 x}{(a_2^2 x^2 + 4c_k^2)^{3/2}} (a_1(1-a_1) - a_2(1-a_2)) < 0.$$

By direct evaluation,  $f(0) > 0$  and  $f(\infty) < 0$ , giving the conclusion. The maximizing value  $x_M$  can be conveniently obtained by a very short FORTRAN program that does interval halving, and Table 1 was so obtained.

Table 1  
Maxima of  $r(x)$

$k$	$x_M$	Maximum
3	.711	1.003
5	.781	1.008
10	.847	1.012
20	.885	1.015
50	.911	1.017
100	.919	1.017
1000	.928	1.018

Suppose  $k$  is fixed and  $M$  is the maximum value of  $r(x)$ , obtained as above. Then

$$(1+x^2/k)f_k(x)p(x, \gamma_{\min}) < \bar{F}_k(x) < (1+x^2/k)f_k(x)p(x, \gamma_{\max}) \quad (2.27)$$

and the absolute error for either approximation is bounded by

$(M-1)(1+x^2/k)f_k(x)p(x, \gamma_{\min})$  and for any given  $x$  is less than

$(1+x^2/k)f_k(x)(p(x, \gamma_{\max}) - p(x, \gamma_{\min}))$ , and the absolute value of the relative error is bounded by  $M-1$  and for any given  $x$  is less than  $p(x, \gamma_{\max})/p(x, \gamma_{\min}) - 1$ .

Some numerical examples are given in Table 2. Here  $U_{\max}$  is the right-hand and  $U_{\min}$  the left-hand expression of (2.27).

Table 2  
Upper and Lower Bounds to t-Tail Probabilities

k	x											
	.1		.5		1.0		2.0		4.0		6.0	
	$U_{\min}$	$U_{\max}$	$U_{\min}$	$U_{\max}$	$U_{\min}$	$U_{\max}$	$U_{\min}$	$U_{\max}$	$U_{\min}$	$U_{\max}$	$U_{\min}$	$U_{\max}$
3	.463	.463	.325	.326	.195	.196	.696-01	.696-01	.140-01	.140-01	.464-02	.464-02
5	.461	.462	.318	.320	.181	.182	.509-01	.512-01	.516-02	.517-02	.923-03	.924-03
10	.460	.461	.312	.315	.169	.171	.366-01	.369-01	.126-02	.126-02	.660-04	.662-04
20	.459	.461	.308	.312	.163	.166	.295-01	.298-01	.352-03	.353-03	.362-05	.365-05
50	.458	.461	.306	.311	.160	.162	.254-01	.257-01	.105-03	.105-03	.109-06	.110-06
100	.458	.460	.306	.310	.158	.161	.240-01	.243-01	.607-04	.611-04	.159-07	.159-07
1000	.458	.460	.305	.310	.157	.160	.228-01	.231-01	.340-04	.342-04	.138-08	.138-08



### 3. An Application to the Calculation of Bonferroni t-Percentiles

Let  $T_1, \dots, T_n$  be statistics associated with an experiment such that under the overall null hypothesis  $H_0$  each  $T_i$  has a t-distribution with  $k$  degrees of freedom,  $i = 1, \dots, n$ . Then if it is desired to control the probability error rate (the probability of 1 or more false statements under  $H_0$ ), one possibility is to reject the null-hypothesis associated with  $T_i$  if

$$|T_i| > t_{\alpha/2n},$$

where  $\bar{F}_k(t_{\alpha/2n}) = \alpha/2n$ , and to accept  $H_0$  if

$$|T_i| \leq t_{\alpha/2n}, \quad 1 \leq i \leq n.$$

Here  $t_{\alpha/2n}$  is called the 100  $\alpha/2n$  Bonferroni t-percentile. Similarly we may define the descriptive Bonferroni level by

$$nP_{H_0}[|T_1| > |t_m|], \quad (3.1)$$

where  $|t_m|$  is the maximum of the observed values of  $|T_1|, \dots, |T_n|$ , and the actual descriptive level,  $P_{H_0}[\text{Max}_{1 \leq i \leq n} |T_i| > |t_m|]$ , will be less than or equal to the descriptive Bonferroni level. Hence it is of interest to find  $t_{\alpha/2n}$  and to evaluate probabilities of the type in (3.1). Recently Bailey (1977) has given a table of  $t_{\alpha/2n}$  for small  $n$  and  $n$  associated with pairwise comparisons. The method to be described below has the advantages of simple computer implementation (short program, no tables to be stored), and of working for any  $\alpha$  and  $n$ .

Denote by  $t_{\max}$  the unique solution in  $t$  of

$$(1+t^2/k)f_k(t)p(t, \gamma_{\max}) = \alpha/2n, \quad (3.2)$$

and by  $t_{\min}$  the unique solution in  $t$  of

$$(1+t^2/k)f_k(t)p(t, \gamma_{\min}) = \alpha/2n . \quad (3.3)$$

Then it immediately follows from (2.27) that

$$t_{\min} < t_{\alpha/2n} < t_{\max} ,$$

and hence either may be used as an approximation, or the digits common to both. The solutions to (3.2) and (3.3) can be obtained very simply by a short FORTRAN program that does interval halving. Table 3 was generated this way. If the entries in Bailey's Table are rounded to three decimals, they all fall within the intervals in Table 3. The maximum difference between the upper and lower bound in Table 3 is .006 and this occurs for  $\alpha = .05$ ,  $n = 1$ . In general, the smaller  $\alpha$  and the larger  $n$ , the better the approximation. The  $n$  and  $\alpha$  for Table 3 were chosen deliberately to correspond to Bailey's Table.

Table 3  
Approximations to Bonferroni t-Percentiles

k	$\alpha$															
	.01								.05							
	n				n				n				n			
	1	20	91	190	1	20	91	190	1	20	91	190	1	20	91	190
	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$	$t_{\min}$	$t_{\max}$
3	5.641	5.842	16.326	16.327	27.131	27.131	34.698	34.698	3.182	3.184	9.465	9.465	15.816	15.817	20.253	20.253
5	4.032	4.034	7.976	7.976	10.962	10.962	12.758	12.758	2.570	2.573	5.604	5.605	7.817	7.817	9.136	9.137
10	3.169	3.172	5.049	5.050	6.139	6.140	6.715	6.716	2.227	2.231	4.004	4.006	4.985	4.986	5.496	5.498
20	2.845	2.848	4.146	4.148	4.796	4.798	5.116	5.118	2.084	2.090	3.455	3.457	4.105	4.107	4.421	4.422
50	2.677	2.680	3.723	3.725	4.199	4.201	4.423	4.424	2.007	2.012	3.184	3.186	3.692	3.694	3.928	3.929
100	2.625	2.629	3.598	3.600	4.027	4.029	4.226	4.227	1.982	1.988	3.101	3.104	3.570	3.572	3.784	3.785
1000	2.580	2.583	3.492	3.494	3.883	3.884	4.061	4.062	1.960	1.966	3.030	3.033	3.466	3.468	3.662	3.664

The probabilities in (3.1) may be bounded by (2.27) and numerical examples have been given in Table 2.

The listings of the three very short computer programs referred to in this paper (maximum, bounds, percentiles) are available on request from the author.

# Appendix: Results for the $c_k$

For ease of reference, all the necessary results involving  $c_k$ 's are given here in the form of lemmas.

Lemma A1: The even and the odd  $c_k$ 's form increasing sequences with limit  $1/(2\pi)^{1/2}$ .

Proof: That the limit is  $1/(2\pi)^{1/2}$  is well-known. By the recursion formula for the  $\Gamma$  function,

$$\frac{c_{2i+3}}{c_{2i+1}} = \left(\frac{i+1}{i+1/2}\right) \left(\frac{2i+1}{2i+3}\right)^{1/2} > 1, \quad i \geq 0.$$

Similarly

$$\frac{c_{2i+2}}{c_{2i}} = \left(\frac{i+1/2}{i}\right) \left(\frac{i}{i+1}\right)^{1/2} > 1, \quad i \geq 1,$$

completing the proof.

Lemma A2: For  $k \geq 1$ ,  $1 - 4c_k^2 > 0$ .

Proof: This is proved differently in Soms (1977), but for the sake of completeness a shorter proof is given here. By Lemma A1, it suffices to show that

$$1/(2\pi) < 1/4,$$

which is true.

Lemma A3: For  $k \geq 3$ ,  $8c_k^2 - 1 > 0$ , and for  $k = 1$ ,  $8c_k^2 - 1 < 0$ .

Proof: By direct computation,  $8c_1^2 - 1 < 0$ . From Lemma A1, it suffices to show that  $8c_3^2 = 16/(\pi\sqrt{3}) > 1$  and  $8c_4^2 = 9/8 > 1$ , which is true.

Lemma A4: For  $k \geq 3$ ,  $4c_k^2/(1-4c_k^2) < k/(2(k+2)c_k^2) - 1$ , and for  $k = 1$  the inequality is reversed.



Proof: For  $k \geq 3$ , the inequality is equivalent to

$$c_k^2 < k/(6k+4), \quad (A1)$$

and for  $k = 1$ , to

$$c_1^2 > 1/10. \quad (A2)$$

(A1) will follow from Lemma A5 and (A2) is verified directly.

Lemma A5: For  $k \geq 3$ ,  $c_k^2 < k/(6k+4)$  and for  $k = 1$ ,  $c_k^2 > k/(6k+4)$ .

Proof: The case  $k = 1$  was shown to hold in Lemma A4. Since  $1/2\pi < 1/(6+4/16)$ ,  $c_k^2 < k/(6k+4)$  for  $k \geq 16$ . It is verified directly that it holds for  $3 \leq k \leq 15$ .

## References

- Bailey, B.J.R. (1977), "Tables of the Bonferroni t Statistic," Journal of the American Statistical Association, 72, 469-78.
- Boyd, A. V. (1959), "Inequalities for Mill's Ratio," Reports of Statistical Applications Research, Japanese Union of Scientists and Engineers, 6, 44-6.
- Johnson, Norman L., and Kotz, Samuel (1970), Continuous Univariate Distributions-2, Boston: Houghton Mifflin Co.
- Soms, Andrew P. (1976), "An Asymptotic Expansion for the Tail Area of the t-Distribution," Journal of the American Statistical Association, 71, 728-30.
- Soms, Andrew P. (1977), "Bounds for the t-Tail Area," to appear as a University of Wisconsin Mathematics Research Center Technical Summary Report.

② Technical summary rept.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1807	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER ⑭ MRL-TSR-1807
4. TITLE (and Subtitle) ⑥ RATIONAL BOUNDS FOR THE t-TAIL AREA WITH AN APPLICATION TO BONFERRONI t-PERCENTILES		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) ⑩ Andrew P./Soms		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) ⑮ DAAG29-75-C-0024
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 4 - Probability, Statistics and Combinatorics
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE ⑪ Nov 1977
		13. NUMBER OF PAGES 18 ⑫ 22p.
		15. SECURITY CLASS. (for this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P.O. Box 12211 Research Triangle Park North Carolina 27709 University of Wisconsin-Milwaukee Graduate School		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bonferroni percentiles, bounds, descriptive Bonferroni level, t-tail area, t-distribution.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The bounds of Boyd (1959) for the normal distribution are extended to the t-distribution. It is shown how these bounds can be used to calculate the Bonferroni descriptive level as well as Bonferroni t-percentiles. The adequacy of the approximation is discussed and numerical examples provided.		

221200